- "You are not in control"
 - You do control the consequences but not the decision of what to do about the substance abuse (SA)
- Stages of Change characteristics & interventions
 - "I don't have a problem..." (pre-contemplative)
 - "I might have a problem..." (contemplative)
 - o "I've got a problem and want to work on it" (Action)
 - o "I used again, what do I do?" (Relapse and recovery)
- Interventions
 - Direct intervention
 - Counseling & AA appears to be the most successful
- Factors affecting prevalence towards SA
 - o Genetics
 - Family & extended family eg grandparents
 - o Community
 - o Family abuse
 - Mental illness (use drugs to cover up symptoms of mental illness) can be very dangerous
- Types of addiction
 - o Periodic
 - o Social Drinker
 - o Binger
 - o Heavy drinker
 - o Out of control
 - o Under-cover drinker
- Jellineck's Curve see http://www.in.gov/judiciary/ijlap/jellinek.pdf
- "The most successful intervention" AA
- Relapse & Recovery
- Rehab characteristics
- Only drugs one can die of in withdrawal
 - o ETOH alcohol
 - o Barbiturates
 - o Benzodiazepines
- Australian study on people 6 weeks after stopping smoking THC marijuana
- Pit-falls of recovery programs
- Addicts are probably the most manipulative and smartest at getting their drug of choice

- while still using with no desire, their drug of choice (DOC) has priority over EVERYTHING in their life.
- Danger's while in black-outs
- Help for the family w/ some liabilities
 - Help family to know they are not alone
- Suicide potential
- Weapon potential
- Hiding DOC's
- CAGE assessment
 - CAGE Questions Adapted to Include Drug Use (CAGE-AID)
 - O 1. Have you ever felt you ought to cut down on your drinking or drug use?
 - O 2. Have people annoyed you by criticizing your drinking or drug use?
 - O 3. Have you felt bad or guilty about your drinking or drug use?
 - 4. Have you ever had a drink or used drugs first thing in the morning to steady your nerves or to get rid of a hangover (eye-opener)?
 - Scoring
 - 0 for a "NO"
 - 1 for "yes"
 - Score of 2 or more is "clinically significant" indicates a high potential for problem w/ SA
- Decisions
- Cycle of Use see attachment